

## Strain relatedness in gram-negative bacteremia: Cause or contamination?

Gram-negative bacteremia

Ilkay Bahceci<sup>1</sup>, Aziz Ramazan Dilek<sup>1</sup>, Kazim Sahin<sup>1</sup>, Ilknur Esen Yildiz<sup>2</sup>, Omer Faruk Duran<sup>3</sup>, Zihni Acar Yazici<sup>1</sup>  
Department of Microbiology, Recep Tayyip Erdoğan University, Medical Faculty Hospital

<sup>2</sup>Department of Infectious Diseases, Recep Tayyip Erdoğan University, Medical Faculty Hospital

<sup>3</sup>Department of Medical Microbiology, Recep Tayyip Erdoğan University Medical Faculty, Rize, Turkey

### Abstract

**Aim:** Bloodstream infections are a major cause of mortality, 25% of which are associated with gram-negative bacteremia. To avoid the inappropriate use of antibiotics, it is important to differentiate the bacteremia from contamination. In general, gram-positive bacteria were more likely to be contaminants than gram-negative-bacteria. There is little information in the literature concerning the epidemiology of gram-negative bacteria isolated from sequential blood cultures. Therefore, we aimed to examine the molecular epidemiology of gram-negative bacteria isolated from sequential blood cultures.

**Material and Methods:** A total of 56 patients (112 samples and strains) with two or more sequential positive blood cultures for gram-negative bacteria with the same antibiogram were included in the study. Pulsed-field gel electrophoresis (PFGE) and arbitrarily primed PCR (AP-PCR) were performed for the determination of strain relatedness.

**Results:** While PFGE analysis demonstrated relatedness in 6 isolates, AP-PCR demonstrated 9 relatedness in 112 isolates.

**Discussion:** The results of our study suggest that, although the possibility of contamination is very low in gram-negative bacteremia, this can still take place, as shown in sequential blood cultures with the same antibiogram.

### Keywords

Bacteremia; Strain relatedness; Pulsed-field gel electrophoresis; Arbitrarily primed PCR

DOI: 10.4328/ACAM.20490 Received: 2021-01-16 Accepted: 2021-03-11 Published Online: 2021-03-28 Printed: 2021-08-01 Ann Clin Anal Med 2021;12(8):900-902

Corresponding Author: Ilkay Bahceci, Recep Tayyip Erdoğan University, Medical Faculty Hospital, Microbiology Department, Zihni Derin Campus, Fener street, 53100 Rize, Turkey.

E-mail: mdilkaybahceci@gmail.com P: +90 464 223 61 26

Corresponding Author ORCID ID: <https://orcid.org/0000-0003-3662-169>

## Introduction

Nosocomial infections contribute to significant morbidity and mortality and occur in approximately 4% of hospitalized patients [1]. Bloodstream infection (BSI) is a major cause of mortality and healthcare costs worldwide [2]. BSI causes the prolonged length of stay in hospital and mortality in many cases. BSI is an acute event leading to life-threatening organ dysfunctions such as sepsis and septic shock [3]. Accurate diagnosis and appropriate antimicrobial treatment are essential in order to increase the survivability of the patient and to decrease high rates of morbidity and mortality [4].

The culturing of blood for pathogens is a simple procedure and provides information essential for the evaluation of a variety of infectious diseases [5]. One-fourth of sepsis cases are associated with gram-negative bacteremia [6]. Gram-negative bacilli such as Enterobacteriaceae and *Pseudomonas aeruginosa* are the leading causes of nosocomial bloodstream infections. Antibiotic-resistant strains have emerged among the gram-negative bacilli and are being increasingly recognized [7]. This significant increase in the incidence of infections caused by antibiotic-resistant Gram-negative bacilli has been a great concern in recent years.

To avoid the inappropriate use of antibiotics, it is important to differentiate the bacteremia from contamination. In general, gram-positive bacteria were more likely to be contaminants than were gram-negative-bacteria [8]. Appropriate antimicrobial therapy has been demonstrated to reduce mortality in Gram-negative bacteremia patients [9]. There is only a little information in the literature about the epidemiology of gram-negative bacteria isolation from sequential blood cultures. The aim of the current study was to examine the molecular epidemiology of gram-negative bacteria isolated from sequential blood cultures.

## Material and Methods

Approval for the study was obtained from the institutional review board (No: 18824186-799). A total of 56 patients (112 samples and strains) with two or more sequential positive blood cultures for gram-negative bacteria that had the same antibiograms within a one-week period were included in the study. Clinical data including age, sex, underlying disease, symptoms of infection, other infection sites, and exposure to antibiotic therapy were collected.


Isolation of gram-negative bacteria from blood cultures was carried out using the Bact/Alert automated system (bioMerieux Industry, Hazelwood, MS, USA). The isolated strains were identified by the VITEK 2 system, using GN cards. All isolates were stored frozen at -70°C. Susceptibility testing for all isolates was performed by a disk-diffusion assay using antibiotic disks (Oxoid, UK) on Mueller–Hinton agar in a rich CO<sub>2</sub> atmosphere and by VITEK 2 GNS cards. Pulsed-field gel electrophoresis (PFGE) of the samples treated with XbaI restriction enzyme (Vivantis) was performed by means of a commercially available kit (GenePath Group 1, Bio-Rad Laboratories, Hercules, CA). Strain relatedness was defined as identical (no difference in banding), closely related (2 to 3 different bands), or unrelated (>3 different bands) [10]. In addition, we used a well-known effective PCR method (arbitrarily primed PCR: AP-PCR) for

genotyping of the strains. To prepare templates for AP-PCR, single colonies were inoculated into L broth and were incubated as stationary cultures at 37°C for about 16 hours. Then, 0.5 ml of the resultant stationary-phase culture was boiled for 10 min, diluted 10-fold in distilled water, and used immediately or stored at 4°C until needed. Phenol-extracted template DNA was prepared for AP-PCR. The previously described arbitrary primers (P1:5'-AAGAG CCGT; P2: 5'-CCGC GCCAA; P3:5'-AACGCGCAAC; P4:5'-GCGATCCCCA) were used [11]. Gels were photographed.

## Results

Demographic data of the patients are shown in Table 1.

A total of 112 episodes of bacteremia were identified. The majority of these episodes were due to *Escherichia coli* (26 patients, 52 isolates), with the remainder due to *Acinetobacter baumannii* (19 patients, 38 isolates), *Klebsiella pneumoniae* (7 patients, 14 isolates) and *Klebsiella oxytoca* (4 patients, 8 isolates). Sequential isolates had similar antibiotic sensitivity. Most of the strains were extended-spectrum  $\beta$ -lactamase positive. PFGE analysis demonstrated relatedness in 4 out of 112 isolates (Figure 1). These isolates were *E. coli*. The remaining isolates were identical. AP-PCR analysis revealed differences in six *E. coli*, two *K. pneumoniae* and one *A. baumannii* isolate.



**Figure 1.** Pulsed field gel electrophoretic patterns of non-identical isolates. The figure shows unrelated sequential isolates of patient 35 (35-1, 35-2). M: Ladder.

**Table 1.** Demographic features of the patients

Mean age	48.28 ± 11.6	
Female / male	32/24	
Antibiotic therapy (yes / no)	51/5	
Source of infection	Lung	22 (%39.2)
	Abdomen	14 (%25)
	Urinary Tract	11 (%19.6)
	Wound	9 (%16)
Underlying disease	Malignancy	23
	Diabetes	19
	Other	14

## Discussion

Understanding epidemiological and microbiological data is critical in order to struggle with potentially life-threatening infections such as BSI, because understanding pathogens and their resistance patterns is of great importance for successful treatment. In this study, molecular epidemiology of gram-negative bacteria isolated from sequential blood cultures was investigated. However, resistance patterns are not discussed in detail here, since they are beyond the scope of this study.

Blood culture is a critical tool for the healthcare professional, a positive blood culture can suggest a definitive diagnosis but, like any test, false positive results can limit the utility of this important tool [12]. Contamination is not uncommon during the sample preparation process, which creates serious problems for interventions. Most contaminants are probably introduced from the patient's skin during blood collection [5]. Frequent contaminants included coagulase-negative staphylococci, Bacillus species and Corynebacterium species [8]. Several clinical studies of bloodstream infections have provided guidelines for differentiating true pathogens from contaminants, but there is no gold standard for differentiating pathogens from contaminants [5].

Gram-negative bacteremia accounts for approximately 25% to 27% of bloodstream infections [13]. Gram-negative bacilli such as Enterobacteriaceae and *P. aeruginosa* are the leading causes of nosocomial bloodstream infections, and appropriate antimicrobial therapy has been shown to reduce mortality among patients with gram-negative bacteremia [7]. The identification of the organism has been shown to be the most important prognostic factor in a predictive model for differentiating contamination from infection in blood culture results of bacteremia [12]. In general, gram-positive bacteria were more likely to be contaminants than gram-negative bacteria [8]. It is accepted that *E. coli* and other members of the Enterobacteriaceae and *P. aeruginosa* always or nearly always (>90%) represent true bacteremia, which our study confirms [14].

One proven methodology that can help differentiate blood culture contamination from a true infection is the number of blood culture sets that grow organisms. For true bacteremia, multiple blood culture sets usually grow the same organism [12]. In our study, we included two or more sequential positive blood cultures for gram-negative bacteria with the same antibiograms within a one-week period. We found 4 related *Escherichia coli* isolates and two closely related isolates of *Klebsiella* (*K. pneumoniae* and *A. baumannii*).

Even though there are numerous relatedness studies on gram-positive bacteria, there are very few studies in the literature on the gram-negatives [15-21]. We believe that this is due to overlooking gram-negative bacteria as potential contaminants. In conclusion, the results of our study suggest that although the possibility of contamination is very low in gram-negative bacteremia, it can still take place even in sequential blood cultures with the same antibiograms. This must be taken into account in bacteremia, especially when antibiotic therapy fails.

### Scientific Responsibility Statement

The authors declare that they are responsible for the article's scientific content including study design, data collection, analysis and interpretation, writing, some

of the main line, or all of the preparation and scientific review of the contents and approval of the final version of the article.

### Animal and human rights statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. No animal or human studies were carried out by the authors for this article.

**Funding:** None

### Conflict of interest

None of the authors received any type of financial support that could be considered potential conflict of interest regarding the manuscript or its submission.

### References

- Colpan A, Akinci E, Erbay A, Balaban N, Bodur H. Evaluation of Risk Factors for Mortality in Intensive Care Units: A Prospective Study from a Referral Hospital in Turkey. *Am J Infect Control*. 2005;33(1):42-7.
- Skogberg K, Lyytikäinen O, Ollgren J, Nuorti JP, Ruutu P. Population-based burden of bloodstream infections in Finland. *Clin Microbiol Infect*. 2012; 18(6):170-6.
- Loonen AJM, de Jager CPC, Tossierams J, Kusters R, Hilbink M, Wever PC, et al. Biomarkers and molecular analysis to improve bloodstream infection diagnostics in an emergency care unit. *PLoS One*. 2014;9(1):e87315.
- Liu V, Escobar GJ, Greene JD, Soule J, Whippy A, Angus DC, et al. Hospital deaths in patients with sepsis from 2 independent cohorts. *JAMA*. 2014;312(1):90-2.
- Lee CC, Lin WJ, Shih HI, Wu CJ, Chen PL, Lee HC, et al. Clinical significance of potential contaminants in blood cultures among patients in a medical center. *J Microbiol Immunol Infect*. 2007; 40(5):438-44.
- Wendt C, Messer SA, Hollis RJ, Pfaller MA, Wenzel RP, Herwaldt LA. Recurrent gram-negative bacteremia: incidence and clinical patterns. *Clin Infect Dis*. 1999; 28:611-7.
- Kang CI, Kim SH, Park WB, Lee KD, Kim HB, Kim EC, et al. Bloodstream infections caused by antibiotic-resistant gram-negative bacilli: risk factors for mortality and impact of inappropriate initial antimicrobial therapy on outcome. *Antimicrob Agents Chemother*. 2005; 49(2):760-6.
- Bryan CS. Clinical implications of positive blood cultures. *Clin Microbiol Rev*. 1989; 2(4):329-53.
- Kang C-I, Kim S-H, Kim H-B, Park S-W, Choi Y-J, Oh M-D, et al. *Pseudomonas aeruginosa* bacteremia: risk factors for mortality and influence of delayed receipt of effective antimicrobial therapy on clinical outcome. *Clin Infect Dis*. 2003; 37(6):745-51.
- Neoh HM, Tan XE, Sapri HF, Tan TL. Pulsed-field gel electrophoresis (PFGE): A review of the "gold standard" for bacteria typing and current alternatives. *Infect Genet Evol*. 2019;74:103935. DOI: 10.1016/j.meegid.2019.103935.
- Madico G, Akopyants NS, Berg DE. Arbitrarily primed PCR DNA fingerprinting of *Escherichia coli* O157:H7 strains by using templates from boiled cultures. *J Clin Microbiol*. 1995;33(6):1534-6.
- Hall KK, Lyman JA. Updated review of blood culture contamination. *Clin Microbiol Rev*. 2006; 19:788-802.
- Marschall J, Doherty J, Warren DK. The epidemiology of recurrent Gram-negative bacteremia in a tertiary-care hospital. *Diagn Microbiol Infect Dis*. 2010; 66:456-9.
- Weinstein MP. Blood culture contamination: persisting problems and partial progress. *J Clin Microbiol*. 2003; 41(6):2275-8.
- Khatib R, Riederer KM, Clark JA, Khatib S, Briski LE, Wilson FM. Coagulase-negative staphylococci in multiple blood cultures: strain relatedness and determinants of same-strain bacteremia. *J Clin Microbiol*. 1995; 33(4):816-20.
- Wendt C, Messer SA, Hollis RJ, Pfaller MA, Wenzel RP, Herwaldt LA. Molecular epidemiology of gram-negative bacteremia. *Clin Infect Dis*. 1999; 28:605-10.
- Liao CH, Lai CC, Chen SY, Huang YT, Hsueh PR. Strain relatedness of methicillin-resistant *Staphylococcus aureus* isolates recovered from patients with repeated bacteremia. *Clin Microbiol Infect*. 2010; 16:463-9.
- Cheng AC, Murdoch DR, Harrell LJ, Barth Reller L. Clinical profile and strain relatedness of recurrent enterococcal bacteremia. *Scand J Infect Dis*. 2005; 37:642-6.
- Al Wohoush I, Rivera J, Cairo J, Hachem R, Raad I. Comparing clinical and microbiological methods for the diagnosis of true bacteremia among patients with multiple blood cultures positive for coagulase-negative staphylococci. *Clin Microbiol Infect*. 2011; 17(4):569-71.
- Zaidi AK, Harrell LJ, Rost JR, Reller LB. Assessment of similarity among coagulase-negative staphylococci from sequential blood cultures of neonates and children by pulsed-field gel electrophoresis. *J Infect Dis*. 1996; 174(5):1010-4.
- Harrington SM, Ross TL, Gebo KA, MerzWG. Vancomycin resistance, esp, and strain relatedness: a 1-year study of enterococcal bacteremia. *J Clin Microbiol*. 2004; 42(12):5895-8.

### How to cite this article:

Ilkay Bahceci, Aziz Ramazan Dilek, Kazim Sahin, Ilknur Esen Yildiz, Omer Faruk Duran, Zihni Acar Yazıcı. Strain relatedness in gram-negative bacteremia: Cause or contamination? *Ann Clin Anal Med* 2021;12(8):900-902